



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MBBS

Doctors Data Inc

123 Main St.

St. Charles, 60174 USA

Patient: Sample Patient

Id: 999999

Age: 70 DOB: 12/20/1951

Sex: Female

Sample Collection

Date/Time

Date Collected

07/04/2022

Date Received

07/11/2022

Date Reported


07/18/2022

### Microbiome Abundance and Diversity Summary

The abundance and diversity of gastrointestinal bacteria provide an indication of gastrointestinal health, and gut microbial imbalances can contribute to dysbiosis and other chronic disease states. The GI360™ Microbiome Profile is a gut microbiota DNA analysis tool that identifies and characterizes more than 45 targeted analytes across six Phyla using PCR and compares the patient results to a characterized normobiotic reference population. The web chart illustrates the degree to which an individual's microbiome profile deviates from normobiosis.

**LEGEND**

The web image shows the relative diversity and balance among bacteria belonging to the six primary Phyla. The white shaded area represents the patient's results compared to a normobiotic reference population. The center of the web represents less abundance while the outer edges represent more than normobiotic.




### Dysbiosis and Diversity Index

These indexes are calculated from the results of the Microbiome Profile, with scores ranging from 1 to 5, and do not include consideration of dysbiotic and pathogenic bacteria, yeast, parasites and viruses that may be reported in subsequent sections of the GI360™ test.

The Dysbiosis Index the (DI) is calculated strictly from the results of the Microbiome Profile, with scores from 1 to 5. A DI score above 2 indicates dysbiosis; a microbiota profile that differs from the defined normobiotic reference population. The higher the DI above 2, the more the sample deviates from the normobiotic profile. The dysbiosis test and DI does not include consideration of dysbiotic and pathogenic bacteria, yeast, parasites and viruses that may be reported in subsequent sections of the GI360™ test.

A diversity score of 3 indicates an expected amount of diversity, with 4 & 5 indicating an increased distribution of bacteria based on the number of different species and their abundance in the sample, calculated based on Shannon's diversity index. Scores of 1 or 2 indicate less diversity than the defined normobiotic reference population.



Dysbiosis Index



Diversity Score



### Key Findings

- Butyrate producing bacteria
- Gut barrier protective bacteria
- Gut intestinal health marker
- Pro-inflammatory bacteria
- Gut barrier protective bacteria vs. opportunistic bacteria

= Expected     = Imbalanced



# Microbiome Bacterial Abundance; Multiplex PCR



**Order:** 999999-9999



**Client #:** 999999

**Doctor:** Sample Doctor, MBBS  
Doctors Data Inc  
123 Main St.  
St. Charles, 60174 USA

**Patient:** Sample Patient

**Id:** 999999

**Age:** 70 **DOB:** 12/20/1951

**Sex:** Female

**Sample Collection**

**Date Collected**

**Date Received**

**Date Reported**


**Date/Time**

07/04/2022


07/11/2022

07/18/2022

## LEGEND



Results are graphed as deviations from a normobiotic population. Normobiosis or a normobiotic state characterizes a composition of the microbiota profile in which microorganisms with potential health benefits predominate in abundance and diversity over potentially harmful ones.



### Notes:

The gray-shaded area of the bar graph represents reference values outside the reporting limits for this test.

\*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.

Methodology: Multiplex PCR



# Microbiome Bacterial Abundance; Multiplex PCR



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MBBS  
Doctors Data Inc  
123 Main St.  
St. Charles, 60174 USA

Patient: Sample Patient

Id: 999999

Age: 70 DOB: 12/20/1951

Sex: Female

Sample Collection

Date Collected

Date Received


Date Reported

Date/Time

07/04/2022

07/11/2022

07/18/2022



### Notes:

The gray-shaded area of the bar graph represents reference values outside the reporting limits for this test.

\*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.

Methodology: Multiplex PCR



**Order:** 999999-9999



**Client #:** 999999

**Doctor:** Sample Doctor, MBBS  
Doctors Data Inc  
123 Main St.  
St. Charles, 60174 USA

**Patient:** Sample Patient

**Id:** 999999

**Age:** 70 **DOB:** 12/20/1951

**Sex:** Female

**Sample Collection**

**Date Collected**

**Date Received**

**Date Reported**

**Date/Time**

07/04/2022

07/11/2022


07/18/2022

Firmicutes	Result	-3	-2	-1	0	+1	+2	+3	Reference Interval
<i>Streptococcus salivarius</i> ssp. <i>thermophilus</i>	0				▲				0
<i>Streptococcus</i> spp.	0				▲				0
<i>Veillonella</i> spp.	0				▲				0
Proteobacteria	Result	-3	-2	-1	0	+1	+2	+3	Reference Interval
Proteobacteria	+1					▲			0
<i>Enterobacteriaceae</i>	0				▲				0
<i>Escherichia</i> spp.	0				▲				0
<i>Acinetobacter junii</i>	0				▲				0
Tenericutes	Result	-3	-2	-1	0	+1	+2	+3	Reference Interval
<i>Mycoplasma hominis</i>	0				▲				0
Verrucomicrobia	Result	-3	-2	-1	0	+1	+2	+3	Reference Interval
<i>Akkermansia muciniphila</i>	0				▲				0



## Microbiome Abundance Information:

- The GI360™ Microbiome Profile is a focused gut microbiota DNA analysis tool that identifies more than 45 targeted analytes across six phyla using a CE-marked multiplex PCR system. Patient results are compared to a highly defined normobiotic reference population (n > 1,100). The white shadowed web plot within the hexagonal diagram illustrates the degree to which an individual's microbiome profile deviates from normobiosis. The center of the diagram represents less bacterial abundance while the outer edges represent greater than normobiosis. Deviation from a hexagon-shaped plot indicates variant diversity of the microbial community. Key findings for patient's microbiome profile are summarized in the table below the diagram, and detailed results for all of the analytes are presented on the next 3 pages of the report. Detailed results for the specific bacteria are reported as -3 to +3 standard deviations, as compared to the normobiotic reference population.



### Notes:

The gray-shaded area of the bar graph represents reference values outside the reporting limits for this test.

\*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.

Methodology: Multiplex PCR

**Order:** 999999-9999**Client #:** 999999**Doctor:** Sample Doctor, MBBS

Doctors Data Inc

123 Main St.

St. Charles, 60174 USA

**Patient:** Sample Patient**Id:** 999999**Age:** 70 **DOB:** 12/20/1951**Sex:** Female**Sample Collection****Date/Time****Date Collected**

07/04/2022

**Date Received**

07/11/2022

**Date Reported**

07/18/2022

## Introduction

This analysis of the stool specimen provides fundamental information about the overall gastrointestinal health of the patient. When abnormal microflora or significant aberrations in intestinal health markers are detected, specific commentaries are presented. If no significant abnormalities are found, commentaries are not presented.

## Microbiome Abundance Information

### Actinobacteria (phylum)

Actinobacteria is one of the largest bacterial phyla, comprised of Gram-positive bacteria. This phylum includes a wide range of species, with different morphological and physiological characteristics. Significant groups in the human colon include Actinomycetales and Bifidobacteriales. Actinomycetales were inversely associated with clinically significant depression in IBS patients, suggesting these bacteria may be depleted in depressed IBS patients. A strict vegetarian diet may increase the total count of *Actinomyces* spp. compared to following a Western diet.

### Bacteroidetes (phylum)

Bacteroidetes make up approximately 28% of the gut microbiota in healthy human adults. They are early colonizers of the infant gut and are amongst the most stable, at a species and strain level, in the host. A low preponderance of Bacteroidetes in relation to Firmicutes has been associated with obesity, though this can increase with weight loss and restricted calorie intake.

#### ↑ *Bacteroides* (species)

Species in the genus *Bacteroides* carry out broad metabolic functions, including degradation of complex plant polysaccharides, proteolytic activities, de-conjugation of bile acids, mucosal barrier integrity, short chain fatty acid production, fatty acid storage and glucose metabolism. *Bacteroides* spp. are maintained at a higher abundance in breastfed individuals into adulthood. *Bacteroides fragilis* plays an important role in the prevention of intestinal inflammation. An energy-restricted diet has been shown to increase *B. fragilis* in overweight adolescents. An increase in *B. stercoris* has been associated with higher risk of colon cancer. Decreased levels of *Bacteroides* spp. have been reported in association with multiple sclerosis, rheumatoid arthritis and Parkinson's disease.

#### ↑ *Parabacteroides* (genus)

The abundance of *Parabacteroides* spp., major anaerobic producers of acetate and succinate is increased with a high fat diet and is positively correlated with body weight. *Parabacteroides* spp., along with certain *Bacteroides* spp., have been shown to distinguish healthy adults from patients with irritable bowel syndrome or ulcerative colitis. Reduced abundance of this group of bacteria has also been linked to Crohn's disease in children. *Parabacteroides* spp. has been found to be less abundant in patients with multiple sclerosis.

### Firmicutes (phylum)

The phylum Firmicutes constitutes the most diverse and abundant group of gastrointestinal microbiota which are grouped into four classes, Bacilli, Clostridia, Erysipelotrichia, and Negativicutes. They constitute about 39% of gut bacteria in healthy adults, but may increase to as high as 80% in an imbalanced microbial community.

#### ↑ *Clostridium methylpentosum* (species)

Appropriate digestion and metabolism of complex dietary carbohydrates from plants drives healthy diversity in the gut microbiota. *Clostridium methylpentosum* ferments the naturally occurring sugar L-rhamnose that is released by microbial breakdown of plant-derived pectin. Rhamnose is fermented to propionate and acetate which are short chain fatty acids that have pivotal regulatory roles in the maintenance of mucosal barrier integrity, gut microbiota balance, post-prandial appetite suppression and normoglycemia. Lower levels of *C. methylpentosum* were reported for children with autism and pervasive developmental disorder compared to neurotypicals controls. Consumption of probiotic yogurt LKM512 containing *Bifidobacterium animalis* (subspecies lactis LKM512) increased the levels of *C. methylpentosum*.

#### ↓ *Eubacterium hallii* (species)

*Eubacterium hallii* and *Eubacterium rectale* are both part of the *Lachnospiraceae* family that is in the Firmicutes phylum. *E. hallii* and *E. rectale* produce butyrate that is a key regulator of mucosal barrier integrity and function. Decreased levels of *Eubacterium* spp. have been associated with very high protein diets. *Eubacterium hallii* is capable of metabolizing glucose into products with antimicrobial properties.



**Order:** 999999-9999



**Client #:** 999999

**Doctor:** Sample Doctor, MBBS

Doctors Data Inc

123 Main St.

St. Charles, 60174 USA

**Patient:** Sample Patient

**Id:** 999999

**Age:** 70 **DOB:** 12/20/1951

**Sex:** Female

**Sample Collection**

**Date Collected**

**Date Received**

**Date Reported**

**Date/Time**

07/04/2022

07/11/2022

07/18/2022

**Microbiome Abundance Information continued...**

↓ **Eubacterium rectale (species)**

*Eubacterium rectale* is part of the *Lachnospiraceae* family and produces butyrate. *Eubacterium rectale* was found to be in lower abundance in patients with type 2 diabetes, colorectal cancer, and chronic idiopathic diarrhea. There is a negative correlation between *Eubacterium rectale* levels and the symptomatology of irritable bowel syndrome (IBS). Decreased levels of *Eubacterium* spp. have been associated with very high protein diets.

↑ **Ruminococcus (genus)**

Members of *Ruminococcus* sensu produce acetate, but not butyrate. *Ruminococcus gnavus*, like *Akkermansia muciniphila* is a mucin degrading specialist. Higher levels of *Ruminococcus* spp. were associated with non-alcoholic fatty liver disease and non-alcoholic steatohepatitis. Reduced levels of *R. bromii* were observed in patients with primary biliary cirrhosis. Increased abundance of *Ruminococcus* spp. has been reported in irritable bowel syndrome (IBS), whereas *Ruminococcus* spp. are reportedly decreased in abundance with Crohn's disease and ulcerative colitis. *Ruminococcus gnavus* has been found to be in higher abundance in diarrhea predominant IBS. Intake of resistant starch has been associated with increased levels of *R. bromii*, whereas a diet rich in animal protein and fat was found to reduce the abundance of this species in human gut.

**Proteobacteria (phylum)**

Proteobacteria include a wide variety of pathogens, including species within the *Escherichia*, *Shigella*, *Salmonella*, *Vibrio*, and *Helicobacter* genera. The phylum includes a number of species that are permanent residents of the microbiota and capable of inducing nonspecific inflammation and diarrhea when their presence is increased. Proteobacteria make up approximately 2% of the gut microbiota in healthy adults.

↑ **Proteobacteria**

A high-fat diet is positively associated with an abundance of Proteobacteria. Slightly increased abundance of Proteobacteria may be associated with low-grade inflammation. Proteobacteria are increased in inflammatory bowel disease and irritable bowel syndrome. Higher abundance of Proteobacteria has been associated with a moderate to severe disease course in newly discovered ulcerative colitis patients. They are associated with diarrhea in IBS.

**Tenericutes (phylum)**

Tenericutes are cell wall-less bacteria that do not synthesize precursors of peptidoglycan. Tenericutes consist of four main clades designated as the *Acholeplasma*, *Spiroplasma*, *Pneumoniae* and *Hominis* clusters. Tenericutes are typically parasites or commensals of eukaryotic hosts.

**Verrucomicrobia (phylum)**

Verrucomicrobia is a less common phylum in the human gut microbiota, but one with increasing recognition with regards to health. Verrucomicrobia includes *Akkermansia muciniphila*. The obligate anaerobe *A. muciniphila* constitutes 3-5% of total bacteria in a healthy microbiome, and has a protective or anti-inflammatory role in the intestinal mucosa.