

**Provider:** sample  
**Patient:** sample  
**Accession #:**

**Sex:**  
**Age:**  
**Sample Type:** Urine Card

**Collected:**  
**Received:**  
**Completed:**

Analyte	Result ( $\mu\text{g}/\text{mg}$ creatinine)	Reference Range	Population Percentile	
<b>Glycolysis</b>				
1. Pyruvate (H)	3.16	< 1.90	92%	
2. Lactate	5.87	< 23.35	39%	
<b>Citric Acid Cycle</b>				
3. Citrate	56.15	71.30 - 772.63	6%	
4. Cis-Aconitate	37.40	< 40.54	61%	
5. Isocitrate	39.29	19.94 - 74.88	19%	
6. Alpha-Ketoglutarate (H)	33.73	< 18.94	93%	
7. Succinate	5.34	< 20.99	38%	
8. Fumarate	0.51	< 1.13	61%	
9. Malate	1.04	< 2.62	31%	
<b>Fatty Acid Oxidation</b>				
10. Adipate (H)	6.01	< 4.42	93%	
11. Suberate	1.22	< 2.64	66%	
12. Ethylmalonate	1.87	< 3.88	35%	
13. Methylsuccinate	2.58	< 2.20	85%	
<b>Markers for Protein Metabolism</b>				
14. Alpha-Ketoisovalerate (H)	0.40	< 0.49	90%	
15. Alpha-Ketoisocaproate	<LLOQ	< 1.09	N/A	
16. Alpha-Keto-Beta-Methylvalerate	0.34	< 1.29	23%	
17. Beta-Hydroxyisovalerate	2.37	< 8.86	8%	
18. Methylmalonate	<LLOQ	< 1.64	N/A	
19. Hydroxymethylglutarate	5.66	< 7.20	89%	

Reference range updated 5/21/2021. Reference range is not gender adjusted. Reference range is age adjusted for children. Method: LC/MS/MS. LLOQ: Lower limit of quantitation ULOQ: Upper limit of quantitation. Lactate is reported as D- and L-Lactate combined on OAP. This test is not intended to diagnose, treat, cure, or prevent any disease or replace the medical advice and/or treatment obtained from a qualified healthcare practitioner. US BioTek Laboratories has developed and determined the performance characteristic of this test under the Clinical Laboratory Improvement Amendments (CLIA). This test has not been evaluated by the U.S. Food and Drug Administration. This test does not assess for neonatal inborn errors of metabolism and is based on stable renal function and normal renal clearance.

**Provider:** sample  
**Patient:** sample  
**Accession #:**

**Sex:**  
**Age:**  
**Sample Type:** Urine Card

**Collected:**  
**Received:**  
**Completed:**

Analyte	Result ( $\mu\text{g}/\text{mg}$ creatinine)	Reference Range	Population Percentile	
<b>Ketone Metabolites</b>				
20. Alpha-Hydroxybutyrate	0.21	< 1.24	25%	
21. Beta-Hydroxybutyrate	0.91	< 8.09	54%	
<b>Markers of Neurotransmitter Metabolism</b>				
22. Vanilmandelate	3.26	< 3.64	63%	
23. Homovanillate (H)	8.12	< 6.66	90%	
24. 5-Hydroxyindoleacetate	4.69	1.17 - 8.06	81%	
25. Quinolinate	3.03	< 5.37	28%	
26. Kynurenate	1.88	< 2.49	59%	
<b>Markers of Detoxification</b>				
27. Para-Hydroxyphenyllactate	0.68	< 1.55	81%	
28. Orotate	<LLOQ	< 1.04	N/A	
29. Pyroglutamate	38.45	14.58 - 37.47	90%	
30. Benzoate	<LLOQ	< 6.87	N/A	
31. Hippurate (H)	1101.08	17.13 - 768.53	99%	
<b>Markers of Bacterial Metabolism</b>				
32. Para-Hydroxybenzoate	<LLOQ	< 1.43	N/A	
33. Para-Hydroxyphenylacetate (H)	20.54	< 26.39	90%	
34. 2-Hydroxyphenylacetate	1.16	< 1.24	81%	
35. 3-Indoleacetate (L)	<LLOQ	0.46 - 9.21	N/A	
36. Tricarballylate (H)	1.56	< 1.06	91%	

Reference range updated 5/21/2021. Reference range is not gender adjusted. Reference range is age adjusted for children. Method: LC/MS/MS. LLOQ: Lower limit of quantitation ULOQ: Upper limit of quantitation. Lactate is reported as D- and L-Lactate combined on OAP. This test is not intended to diagnose, treat, cure, or prevent any disease or replace the medical advice and/or treatment obtained from a qualified healthcare practitioner. US BioTek Laboratories has developed and determined the performance characteristic of this test under the Clinical Laboratory Improvement Amendments (CLIA). This test has not been evaluated by the U.S. Food and Drug Administration. This test does not assess for neonatal inborn errors of metabolism and is based on stable renal function and normal renal clearance.