

Correct collection process is vital to receiving the most accurate NGS diagnostic results. Please follow the steps below.



BV/VAGINAL SAMPLE COLLECTION METHOD | FOR WOMEN

THIS TEST COMES WITH:



2 Vaginal Swabs



1 Bio Bag



1 Lab Requisition



Box with
Return Label

COLLECTION DO'S & DON'TS

✓ **DO** - Clean the genital area with soap and water before collecting the sample.

✓ **DO** - Perform a preparatory urination two hours in advance of first sample.

✗ **DO NOT** - Use vaginal suppositories for 24 hours prior to collection.

✗ **DO NOT** - Take antibiotics. Patient must be off all antibiotics for 2 days prior to collection. This is optimal. However, if not possible, the test can still be run.

✗ **DO NOT** - Contaminate the sample with period blood during menstruation.

SWAB METHOD


STEP 1

Clean the genital area with soap and water before collecting the sample.




STEP 2

Twist the cap at the dotted line to break the seal. Remove the Swab completely from the tube not letting your hands touch the swab beyond the break point.



STEP 3

Gently collect the specimen at the vaginal opening/center of the infected area in a circular motion starting near the center and working outwards. Move the swab over as much of the surface of the vaginal area as possible. **Be sure to repeat the process with the second swab.**



STEP 4

Put **BOTH** swabs in the lab bag and tightly seal.



SEE PATIENT INFORMATION &
SHIPPING INSTRUCTIONS ON REVERSE



Correct collection process is vital to receiving the most accurate NGS diagnostic results. Please follow the steps below.

PATIENT INFORMATION

1. MARK SAMPLE

Use a permanent marker to write patient name, date of birth and sample collection date on Vaginal Swab Tubes.



2. SIGN LAB REQ

Please have the patient sign the lab requisition form accepting financial responsibility. Be sure the physician has also signed the lab requisition form confirming consent.



PACKING SAMPLES FOR SHIPMENT

1. Place the 8mL Vacuum Vial and either the 90mL Collection Cup or Multi-Purpose Swab Tube into the centre/seal-able pocket of Biohazard Lab Bag.
2. Place folded Lab Requisition Form into the short pocket of the Lab Bag.
3. IMPORTANT: Place only one Sample and one Lab Requisition in each Lab Bag.
4. Peel strip off Lab Bag to expose adhesive backing and follow instructions printed on Bag to create a continuous, airtight seal.
5. Place each sealed Lab Bag into the white Shipping Box and apply UPS label to the outside and seal.
6. You can add multiple completed biohazard lab bags into 1 white shipping box.



3 SHIPPING METHODS

1. UK Based ? Use the Pre-Paid Royal Mail Return Label
2. Drop off at any Post Office.
3. Or Hand Deliver to Regenerus Laboratories, Aero 14, Kings Mill Lane, Redhill, Surrey RH1 5JY, United Kingdom.
4. If International email shipping@regeneruslabs.com to arrange for your courier label.
5. International shipping fee is not included within your fee.
6. Make sure to retain your tracking number.



THIS TEST COMES WITH:



1 Urine Collection Cup
w/ 8mL Vacuum Vial



1 Bio Bag



1 Lab Requisition



Box with
Return Label

COLLECTION DO'S & DON'TS

- ✓ **DO** - Clean the genital area with soap and water before collecting the sample.
- ✓ **DO** - Perform a preparatory urination two hours in advance of first sample.
- ✓ **DO** - Be sure to press vial onto needle in a smooth, quick motion to ensure successful transfer a sample liquid.
- ✓ **DO** - Place red seal strip (see step 4-b) over the cap attaching firmly to both sides of the vial after taking the sample.

- ✗ **DO NOT** - Use Lidocaine gel. If collecting urine via catheter, Lidocaine gel can render sample invalid or give false negative test results.
- ✗ **DO NOT** - Take in any fluids two hours in advance of first sample.
- ✗ **DO NOT** - Take antibiotics. Patient must be off all antibiotics for 2 days prior to collection. This is optimal. However, if not possible, the test can still be run.
- ✗ **DO NOT** - Fill 8ml vial more than 1/2 full.
- ✗ **DO NOT** - Remove rubber cap from 8ml sample vial. Vials are shipped vacuum sealed. If pressurized transfer fails, pour liquid into 8mL Vial, cap tightly, seal with red strip.

Scan this code with your mobile device to view a video tutorial of the collection process.



COLLECTING URINE SAMPLE WITH VACUTAINER

STEP 1

Cleanse all areas around the urinary exit with soap and water. With a clean tissue, blot off the urethra without touching any part of genital area with hands.



STEP 2

Spread a clean tissue on a surface close to the toilet bowl. Remove the lid from the urine cup and place it top-side down on the tissue. Place the open urine cup next to the tissue, being careful not to touch the rim of the cup or lid with fingers or body surface at any time.



STEP 3

Hold the empty cup in one hand. Begin urination into toilet, not cup, stop urination partway through then continue urination into cup. Keep urinating into cup until cup is 1/3 to 1/2 full, then withdraw cup. Secure lid tightly.




STEP 4

a. Peel yellow label off lid. **CAUTION: Needle under label - DO NOT push finger through label.** Once label is removed, press the yellow rubber capped end of 8ml vial onto the needle to enable vacuum transfer of urine into the 8ml specimen tube. Draw in at least 4ml (half full) of the sample fluid and remove specimen tube with cap intact.



b. Place red seal strip over 8ml vial cap. Ensure seal strip is firmly attached on both sides of vial to prevent cap from coming off during transit.



SEE PATIENT INFORMATION & SHIPPING INSTRUCTIONS ON REVERSE



Correct collection process is vital to receiving the most accurate NGS diagnostic results. Please follow the steps below.

PATIENT INFORMATION

1. MARK SAMPLE

Use a permanent marker to write patient name, date of birth and sample collection date on 8mL Vacuum Vial.




2. SIGN LAB REQ

Please have the patient sign the lab requisition form accepting financial responsibility. Be sure the physician has also signed the lab requisition form confirming consent.



PACKING SAMPLES FOR SHIPMENT

1. Place the 8mL Vacuum Vial and either the 90mL Collection Cup or Multi-Purpose Swab Tube into the centre/seal-able pocket of Biohazard Lab Bag.
2. Place folded Lab Requisition Form into the short pocket of the Lab Bag.
3. **IMPORTANT:** Place only one Sample and one Lab Requisition in each Lab Bag.
4. Peel strip off Lab Bag to expose adhesive backing and follow instructions printed on Bag to create a continuous, airtight seal.
5. Place each sealed Lab Bag into the white Shipping Box and apply UPS label to the outside and seal.
6. You can add multiple completed biohazard lab bags into 1 white shipping box.



3 SHIPPING METHODS

1. UK Based ? Use the Pre-Paid Royal Mail Return Label
2. Drop off at any Post Office.
3. Or Hand Deliver to Regenerus Laboratories, Aero 14, Kings Mill Lane, Redhill, Surrey RH1 5JY, United Kingdom.
4. If International email shipping@regeneruslabs.com to arrange for your courier label.
5. International shipping fee is not included within your fee.
6. Make sure to retain your tracking number.

